

Problem Set 4

Problem 1. Let $R = k[x, y]$, where k is a field, let $Q = \text{frac}(R)$ be the fraction field of R . We are going to show that the R -module $M = Q/R$ is divisible but not injective.

- a) Show that if $ax + by = 0$ for some $a, b \in R$, we must have $b \in (x)$.
- b) Show that $x \mapsto \frac{1}{y}$ and $y \mapsto 0$ induces a well-defined R -module homomorphism $(x, y) \xrightarrow{f} Q/R$.
- c) Show that M is a divisible R -module, but not injective.

Problem 2. Let R be a domain. Show that if R has a nonzero module M that is both injective and projective, then R must be a field.¹

Problem 3. Let R be a Noetherian ring, M a finitely generated R -module, N an R -module, and W a multiplicatively closed subset of R . Show that there is an isomorphism

$$W^{-1} \text{Hom}_R(M, N) \cong \text{Hom}_{W^{-1}R}(W^{-1}M, W^{-1}N).$$

Clearly indicate where you are using the hypotheses that R is Noetherian and M is finitely generated, as they are necessary.²

Problem 4. Let \mathcal{A} be an abelian category.

- a) Show that $\ker(x \xrightarrow{0} y) = 1_x$, $\text{coker}(x \xrightarrow{0} y) = 1_y$, and $\text{im}(x \xrightarrow{0} y) = 0 \rightarrow y$.
- b) Show that f is a mono if and only if $fg = 0$ implies $g = 0$, and g is an epi if and only if $fg = 0$ implies $f = 0$.
- c) Show that f is a mono if and only if $\ker f = 0$, and g is an epi if and only if $\text{coker } g = 0$.
- d) Show that $0 \rightarrow A \xrightarrow{f} B$ is exact if and only if f is a mono.
- e) Show that $B \xrightarrow{g} C \rightarrow 0$ is exact if and only if g is an epi.

Problem 5. Consider an abelian category. If g is an epi and f is a mono, then $\ker(fg) = \ker g$, $\text{coker}(fg) = \text{coker } f$, and $\text{im}(fg) = \text{im } f = f$.

¹Hint: show that any nonzero R -module homomorphism $M \rightarrow R$ must be surjective, and then show that such a homomorphism must exist.

²Hint: start by noting that the obvious map $W^{-1} \text{Hom}_R(M, N) \rightarrow \text{Hom}_{W^{-1}R}(W^{-1}M, W^{-1}N)$ is natural on M and an isomorphism when $M = R^n$. Then apply appropriate functors to a presentation $R^m \rightarrow R^n \rightarrow M$ for M .

An R -module F is *faithfully flat* if F is flat and $F \otimes_R M \neq 0$ for every nonzero R -module M .

Problem 6. Give an example of a module that is flat but not faithfully flat. Show³ that the following are equivalent:

- a) F is faithfully flat.
- b) F is flat and for every proper ideal I , $IF \neq F$.
- c) F is flat and for every maximal ideal \mathfrak{m} , $\mathfrak{m}F \neq F$.
- d) For every sequence of R -modules $A \xrightarrow{f} B \xrightarrow{g} C$, $A \xrightarrow{f} B \xrightarrow{g} C$ is exact if and only if $F \otimes_R A \xrightarrow{1 \otimes f} F \otimes_R B \xrightarrow{1 \otimes g} F \otimes_R C$ is exact.

Problem 7. Consider the ring $R = \mathbb{Q}[x, y, z, a, b, c]/(xb - ac, yc - bz, xc - az)$, the ideal $I = (x, a)$ in R , $I = (x, a)$, and the 2-generated R -module $M = Rf + Rg$, where the generators f, g satisfy the relations

$$yf - xg = 0 \quad bf - cg = 0 \quad cf - zg = 0.$$

Let $S = \mathbb{Q}[x, y, z]$ and P be the ideal in S defining the curve $\{(t^{13}, t^{42}, t^{73}) \mid t \in \mathbb{Q}\}$.

- a) Find the first 6 steps in the minimal free resolutions for R/I and N over R .
- b) Apply $\text{Hom}_R(-, M)$ to the portion of a minimal free resolution you found for R/I . Is this an exact complex? If not, in what homological degrees do we have non-trivial homology?
- c) Find a minimal free resolution for P over S . Make sure your resolution *is* minimal!

³Hints:

- For c) \implies a), for each R -module $M \neq 0$ consider some nonzero $m \in M$ and $I = \text{ann } m$.
- For a) \implies d), show that $\text{im}(1_F \otimes f) = F \otimes_R \text{im } f$ and $\ker(1_F \otimes f) = F \otimes_R \ker g$, and then consider the short exact sequence $0 \rightarrow \text{im } f \rightarrow \ker g \rightarrow \ker g / \text{im } f \rightarrow 0$.
- For d) \implies a), show that for any R -module $M \neq 0$, the identity map on M induces a nonzero map $F \otimes_R M \rightarrow R \otimes_R M$.