

Homework #8

Problems to hand in on Thursday, March 28, in the beginning of class. Write your answers out carefully, staple pages, and write your name and section number on each page.

- 1) Let S^1 be the subset of \mathbb{C} consisting of complex numbers of absolute value 1; that is

$$S^1 := \{z \in \mathbb{C} \mid |z| = 1\}.$$

- (a) Prove that S^1 is a subgroup of \mathbb{C}^\times .

- (b) Prove that the map

$$S^1 \rightarrow \mathrm{SL}_2(\mathbb{R}) \quad x + iy \mapsto \begin{bmatrix} x & -y \\ y & x \end{bmatrix}$$

is an injective group homomorphism.

- (c) Prove that S^1 is isomorphic to $\mathrm{SO}_2(\mathbb{R})$, the group of 2×2 orthogonal matrices of determinant 1. Use this to give a geometric interpretation of the group S^1 that explains why some call it the “continuous rotation group.”
- (d) For every positive integer n , find an element of order n in S^1 .
- (e) Find an element of infinite order in S^1 .

- 2) Towards the end of the worksheet on group homomorphisms, we encountered the following:

THEOREM: If \mathbb{F} is a finite field, then \mathbb{F}^\times is cyclic.

- (a) Check that 2 is not a generator for \mathbb{Z}_{17}^\times but 3 is a generator for \mathbb{Z}_{17}^\times .
- (b) Verify that $\mathbb{F}_9 = \mathbb{Z}_3[x]/(x^2 + x + 2)$ is a field, and find a generator for \mathbb{F}_9^\times .
- (c) Read Corollary 7.10 on page 200, and use this corollary to prove the THEOREM above.¹
- (d) The THEOREM above only applies to finite fields, but we can sometimes describe multiplicative groups of infinite fields in terms of other groups. Show that $\mathbb{R}^\times \cong \mathbb{R} \times \mathbb{Z}_2$.
- (e) Show that $\mathbb{C}^\times \cong \mathbb{R} \times S^1$.

- 3) Consider the following elements in $GL_2(\mathbb{C})$:

$$\mathbf{1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{i} = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}, \quad \mathbf{j} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad \mathbf{k} = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix},$$

Let Q be the subgroup of $GL_2(\mathbb{C})$ generated by the matrices $\mathbf{i}, \mathbf{j}, \mathbf{k}$. You should verify (but not necessarily turn in a proof) that Q contains the 8 elements $\{\pm \mathbf{1}, \pm \mathbf{i}, \pm \mathbf{j}, \pm \mathbf{k}\}$. You may wish to make a multiplication table for Q to answer the following questions. (You do not need to turn in the multiplication table – although notice you have already written in down in last week’s webwork!)

- (a) Find the complete list of all cyclic subgroups of Q of order 4.
- (b) Find the complete list of all cyclic subgroups of Q of order 2.
- (c) Find the complete list of all noncyclic subgroups of Q of order 4.

¹For a hint, look at the worksheet on group homomorphisms.

- (d) Can Q be generated by two elements? Prove it.
- (e) Is Q_8 isomorphic to D_4 ? Prove or disprove.
- 4) Consider the symmetric group \mathbb{S}_n .
- (a) Show that every element of \mathbb{S}_n is a product of transpositions.²
- (b) Let $\tau \in \mathbb{S}_n$ be a permutation, and (ab) be a transposition. Show that $\tau(ab)\tau^{-1} = (\tau(a)\tau(b))$, the transposition changing $\tau(a)$ and $\tau(b)$.
- (c) Show that $(ij) = (1i)(1j)(1i)$. Conclude that every element of \mathbb{S}_n is the product of transpositions of the form $(1i)$.
- (d) Let σ be the n -cycle $(2 \cdots n)$. Show that $(1i) = \sigma^{i-2}(12)(\sigma^{-1})^{i-2}$. Conclude that $\mathcal{S}_n = \langle (12), (2 \cdots n) \rangle$.

²Hint: One possibility for a quick solution is induction on n . Can you multiply any permutation by a transposition to obtain a permutation that fixes one element?