

## Math 412. Adventure sheet on Quotient Groups

Fix an arbitrary group  $(G, \circ)$ .

DEFINITION: A subgroup  $N$  of  $G$  is **normal** if for all  $g \in G$ , the left and right  $N$ -cosets  $gN$  and  $Ng$  are the *same* subsets of  $G$ .

NOTATION: If  $H \subseteq G$  is any subgroup, then  $G/H$  denotes the set of left cosets of  $H$  in  $G$ . Its elements are *sets* denoted  $gH$  where  $g \in G$ . The cardinality of  $G/H$  is called the **index** of  $H$  in  $G$ .

DEFINITION/THEOREM 8.13: Let  $N$  be a normal subgroup of  $G$ . Then there is a well-defined binary operation on the set  $G/N$  defined as follows:

$$G/N \times G/N \rightarrow G/N \quad g_1N \star g_2N = (g_1 \circ g_2)N$$

making  $G/N$  into a group. We call this the **quotient group** " $G$  modulo  $N$ ".

A. WARMUP: Define the *sign map*:

$$S_n \rightarrow \{\pm 1\} \quad \sigma \mapsto 1 \text{ if } \sigma \text{ is even; } \sigma \mapsto -1 \text{ if } \sigma \text{ is odd.}$$

- (1) Prove that sign map is a group homomorphism.
- (2) Use the sign map to give a different proof that  $A_n$  is a normal subgroup of  $S_n$  for all  $n$ .
- (3) Describe the  $A_n$ -cosets of  $S_n$ . Make a table to describe the quotient group structure  $S_n/A_n$ . What is the identity element?

### Solution.

- (1) By definition, if  $\tau$  is a transposition then  $\tau \mapsto -1$ . Given any element  $\sigma \in S_n$ , if we write  $\sigma$  as a product of transpositions, say  $\sigma = \tau_1 \dots \tau_k$ , then  $\sigma \mapsto (-1)^n$ . Now if  $\sigma' \in S_n$  is a product of  $r$  transpositions,  $\sigma\sigma'$  is a product of  $k+r$  transpositions, and

$$\sigma\sigma' \mapsto (-1)^{k+r} = (-1)^k (-1)^r.$$

- (2) By definition,  $A_n$  is the kernel of the sign map, and we have shown that the kernel of a group homomorphism must be a normal subgroup.
- (3) There are two cosets:  $A_n$  and  $S_n \setminus A_n$ , the last one being the set of odd permutations. The identity element in  $S_n/A_n$  is the coset  $A_n$ , and the group  $S_n/A_n$  is isomorphic to  $\mathbb{Z}_2$ .

B. OPERATIONS ON COSETS: Let  $(G, \circ)$  be a group and let  $N \subseteq G$  be a **normal** subgroup.

- (1) Take arbitrary  $ng \in Ng$ . Prove that there exists  $n' \in N$  such that  $ng = gn'$ .
- (2) Take any  $x \in g_1N$  and any  $y \in g_2N$ . Prove that  $xy \in g_1g_2N$ .
- (3) Define a binary operation  $\star$  on the set  $G/N$  of left  $N$ -cosets as follows:

$$G/N \times G/N \rightarrow G/N \quad g_1N \star g_2N = (g_1 \circ g_2)N.$$

Think through the meaning: the elements of  $G/N$  are *sets* and the operation  $\star$  combines two of these sets into a third set: how? Explain why the binary operation  $\star$  is **well-defined**. Where are you using normality of  $N$ ?

- (4) Prove that the operation  $\star$  in (4) is associative.
- (5) Prove that  $N$  is an identity for the operation  $\star$  in (4).
- (6) Prove that every coset  $gN \in G/N$  has an inverse under the operation  $\star$  in (4).
- (7) Conclude that  $(G/N, \star)$  is a group.

- (8) Does the set of **right cosets** also have a natural group structure? What is it? Does it differ from  $G/N$ ?

**Solution.**

(1) Since  $N$  is normal,  $Ng = gN$ . Given  $ng \in Ng = gN$ , there exists  $n' \in N$  such that  $ng = gn'$ .

(2) There exist some  $n_1, n_2 \in N$  such that  $x = g_1n_1$  and  $y = g_2n_2$ . Then

$$xy = g_1n_1g_2n_2 = g_1(n_1g_2)n_2.$$

We assumed that  $N$  is normal, so  $n_1g_2 \in Ng_2 = g_2N$ . Let  $n \in N$  be such that  $n_1g_2 = g_2n$ . Then

$$xy = g_1(n_1g_2)n_2 = g_1(g_2n)n_1 = (g_1g_2)(n_1n) \in (g_1g_2)N.$$

(3) The problem could be that if we can write a coset in two different ways, say  $g_1N = h_1N$ , then when we multiply by another coset, say  $g_2N$ , then there could be two different possible answers for  $(g_1N) \cdot (g_2N)$ :

- One possible answer is  $(g_1g_2)N$ ;
- another possible answer is  $(h_1g_2)N$ .

We need to check that we really only get one answer for each possible product; so we need to check that  $(g_1g_2)N = (h_1g_2)N$ . This is what we just did in the previous question!

A similar problem arises with the second factor. So to check that our operation really is well-defined, we need to take any  $g_1, h_1, g_2, h_2$  such that  $g_1N = h_1N$  and  $g_2N = h_2N$ , and verify that  $(g_1g_2)N = (h_1h_2)N$ . Again, this is what the previous question says. This is equivalent to proving that  $(g_1g_2)(h_1h_2)^{-1} \in N$ .

(4) Now that we know the operation is well-defined, it is easy to check that properties of the operation on  $G$  pass to  $G/H$ . In particular,  $\star$  is associative because the operation on  $G$  also is associative:

$$(gN \star hN) \star kN = (gh)N \star kN = ((gh)k)N = (g(hk))N = gN \star (hk)N = gN \star (hN \star kN).$$

(5) Given any  $g \in G$ ,

$$gN \star eN = (ge)N = gN = (eg)N = eN \star gN.$$

(6) Let  $g \in G$ . Then

$$g^{-1}N \star gN = (g^{-1}g)N = N = (gg^{-1})N = gN \star g^{-1}N.$$

(7) We have shown that this is a set with an associative operation for which there is an identity and every element has an inverse, so this is a group.

**C. EASY EXAMPLES OF QUOTIENT GROUPS:**

- (1) In  $(\mathbb{Z}, +)$ , explain why  $n\mathbb{Z}$  is a normal subgroup and describe the corresponding quotient group.
- (2) For any group  $G$ , explain why  $G$  is a normal subgroup of itself. What is the quotient  $G/G$ ?
- (3) For any group  $G$ , explain why  $\{e\}$  is a normal subgroup of  $G$ . What is the quotient  $G/\{e\}$ ?

**Solution.**

(1) We have shown that every subgroup of an abelian group is normal, so  $n\mathbb{Z}$  is a normal subgroup of  $\mathbb{Z}$ . The quotient group is the group  $(\mathbb{Z}_n, +)$ .

(2) For every  $g \in G$ ,  $gGg^{-1} \subseteq G$ , because  $G$  is closed for products. This means that  $G$  is a normal subgroup of  $G$ . The quotient  $G/G$  is the trivial group (with one element).

(3) For every  $g \in G$ ,  $g\{e\} = \{g\} = \{e\}g$ , so the trivial subgroup is normal. The quotient group  $G/\{e\}$  is isomorphic to  $G$ .

D. ANOTHER EXAMPLE. Let  $G = \mathbb{Z}_{25}^\times$ . Let  $N$  be the subgroup generated by  $[7]$ .

- (1) Give a one-line proof that  $N$  is normal.
- (2) List out the elements of  $G$  and of  $N$ . Compute the order of both. Compute the index of  $N$  in  $G$ .
- (3) List out the elements of  $G/N$ ; don't forget that each one is a *coset* (in particular, a set whose elements you should list).
- (4) Give each coset in  $G/N$  a reasonable name. Now make a multiplication table for the group  $G/N$ , using these names. Is  $G/N$  abelian?

**Solution.**

- (1)  $G$  is abelian, so  $N$  is a normal subgroup.
- (2)  $G = \{1, 2, 3, 4, 6, 7, 8, 9, 11, 12, 13, 14, 16, 17, 18, 19, 21, 22, 23, 24\}$  and  $N = \{1, 7, 24, 18\}$ .  
So  $|\mathbb{Z}_{25}^\times| = 5^2 - \frac{25}{5} = 20$  and  $|\langle 7 \rangle| = 4$ . By Lagrange's Theorem  $[\mathbb{Z}_{25}^\times : \langle 7 \rangle] = \frac{20}{4} = 5$ .
- (3)  $N = \{1, 7, 24, 18\}$ ,  $2N = \{2, 14, 23, 11\}$ ,  $3N = \{3, 21, 22, 4\}$ ,  $6N = \{6, 17, 19, 8\}$ ,  $9N = \{9, 13, 16, 12\}$ .
- (4) Actually, this is just  $\mathbb{Z}_5$ : it is a group of order 5. So yes, this is an abelian group, and writing a multiplication table is quite easy. What if we wanted to give an explicit isomorphism to  $\mathbb{Z}_5$ ? Our isomorphism must send  $N$  to  $[0]_5$ . Now which element gets sent to  $[1]_5$  does not matter: every element in  $\mathbb{Z}_5$  is a generator! But once we pick what element goes to  $[1]_5$ , the others are completely determined. For example, we can have  $2N \mapsto [1]_5$ ,  $3N \mapsto [2]_5$ ,  $6N \mapsto [4]_5$  and  $9N \mapsto [3]_5$ .

E. THE CANONICAL QUOTIENT MAP: Prove that the map

$$G \rightarrow G/N \quad g \mapsto gN$$

is a group homomorphism. What is its kernel?

**Solution.** Write  $\phi$  for the canonical map. Given  $g, h \in G$ ,  $\phi(gh) = (gh)N = gN \star hN = \phi(g)\phi(h)$ . The kernel of the canonical map is  $N$ . This shows that given any normal subgroup  $N$ , there is always a group homomorphism with kernel  $N$ .

F. INDEX TWO. Suppose that  $H$  is an index two subgroup of  $G$ . Last time, we proved the

THEOREM: *Every subgroup of index two in  $G$  is normal.*

- (1) Describe the quotient group  $G/H$ . What are its elements? What is the table?
- (2) Find an example of an index two subgroup of  $D_n$  and describe its two cosets explicitly. Make a table for this group and describe the canonical quotient map  $G \rightarrow G/H$  explicitly.

**Solution.**

- (1) This is a group of order 2, so isomorphic to  $\mathbb{Z}_2$ . The elements are  $H$  and  $G \setminus H$ .
- (2) The group of rotations! It has  $n$  elements, the  $n$  rotations.

G. PRODUCTS AND QUOTIENT GROUPS: Let  $K$  and  $H$  be arbitrary groups and let  $G = K \times H$ .

- (1) Find a natural homomorphism  $G \rightarrow H$  whose kernel  $K'$  is  $K \times e_H$ .
- (2) Prove that  $K'$  is a normal subgroup of  $G$ , whose cosets are all of the form  $K \times h$  for  $h \in H$ .
- (3) Prove that  $G/K'$  is isomorphic to  $H$ .

**Solution.**

- (1) Consider the projection onto the second component, meaning the map  $\phi : G \rightarrow H$  given by  $\phi(k, h) = h$ . Then  $(k, h) \in K'$  if and only if  $h = e_H$ , or equivalently,  $(k, h) \in K \times e_H$ .
- (2) Since  $K'$  is the kernel of a group homomorphism,  $K'$  is normal. Now note that for each  $h \in H$ ,

$$K \times h = \{(k, h) : k \in K\} = (K \times e_H)(e_G, h).$$

On the other hand, given any  $K$ -coset  $K'(k, h)$ ,  $(e_G, h) = (k^{-1}, e_H)(k, h) \in K'(k, h)$ . So every coset is of the form  $K \times h$  for some  $h$ . Finally, if  $h, h' \in H$ , then  $K \times h = K \times h'$  if and only if  $(e, h')(e, h^{-1}) \in K \times \{e\}$ , or equivalently,  $h'h^{-1} = e$ .

H. What goes wrong if we try to define a group structure on the set of right cosets  $G/H$  where  $H$  is a *non-normal* subgroup of  $G$ ? Try illustrating the problem with the non-normal subgroup  $\langle(1\ 2)\rangle$  in  $S_3$ .

**Solution.**

I. THE FIRST ISOMORPHISM THEOREM. Conjecture and prove **first isomorphism theorem** for groups.

**Solution.** The First Isomorphism Theorem says the following:

Given a surjective group homomorphism  $\phi : G \rightarrow H$ ,  $H \cong G/\ker(\phi)$ .

Here is a proof:

Consider the map  $\psi : G/\ker(\phi) \rightarrow H$  given by  $\psi(\ker(\phi)g) = \phi(g)$ .

This map  $\psi$  is well-defined: given  $g, h \in G$  such that  $\ker(\phi)g = \ker(\phi)h$ , by definition we have  $gh^{-1} \in \ker(\phi)$ , so  $\phi(gh^{-1}) = e$ , and thus  $\phi(g) = \phi(h)$ .

Moreover, this map  $\psi$  is a group homomorphism:

$$\psi(\ker(\phi)(gh)) = \phi(gh) = \phi(g)\phi(h) = \psi(\ker(\phi)g)\psi(\ker(\phi)h).$$