

## Problem Set 4

Due Friday, December 1

Turn in **5** of the following problems. Slightly more challenging problems are indicated by  $(\star)$ .

**Problem 1.** Let  $R = k[x, y]$ , where  $k$  is a field, let  $Q = \text{frac}(R)$  be the fraction field of  $R$ . We are going to show that the  $R$ -module  $M = Q/R$  is divisible but not injective.

- a) Show<sup>1</sup> that if  $ax + by = 0$  for some  $a, b \in R$ , we must have  $b \in (x)$ .
- b) Show that  $x \mapsto \frac{1}{y} + R$  and  $y \mapsto 0$  induces a well-defined  $R$ -module homomorphism  $(x, y) \xrightarrow{f} Q/R$ .
- c) Show that  $M$  is a divisible  $R$ -module, but not injective.

**Problem 2.**  $(\star)$  Let  $R$  be a domain. Show that if  $R$  has a nonzero module  $M$  that is both injective and projective, then  $R$  must be a field.<sup>2</sup>

An  $R$ -module  $F$  is *faithfully flat* if  $F$  is flat and  $F \otimes_R M \neq 0$  for every nonzero  $R$ -module  $M$ .

**Problem 3.**  $(\star)$  Let  $R$  be a commutative ring. Show that the following are equivalent:

- a)  $F$  is faithfully flat.
- b)  $F$  is flat and for every proper ideal  $I$ ,  $IF \neq F$ .
- c)  $F$  is flat and for every maximal ideal  $\mathfrak{m}$ ,  $\mathfrak{m}F \neq F$ .
- d) The complex

$$A \xrightarrow{f} B \xrightarrow{g} C$$

is exact if and only if

$$F \otimes_R A \xrightarrow{1 \otimes f} F \otimes_R B \xrightarrow{1 \otimes g} F \otimes_R C$$

is exact.

**Problem 4.** Let  $M$  be an  $R$ -module. Show that  $M$  is flat if and only if  $\text{Tor}_1^R(M, N) = 0$  for every  $R$ -module  $N$ .

**Problem 5.** Let  $R$  be a commutative ring and  $M$  and  $N$  be  $R$ -modules. Consider the  $R$ -module homomorphism  $f: M \rightarrow M$  given by multiplication by a fixed element  $r \in R$ . Show that the map  $\text{Ext}^i(f, N): \text{Ext}_R^i(M, N) \rightarrow \text{Ext}_R^i(M, N)$  induced by  $f$  is multiplication by  $r$  on  $\text{Ext}_R^i(M, N)$ .

<sup>1</sup>If you know about regular sequences, this is easy to justify. But we aren't assuming anyone has seen regular sequences, so the challenge here is to give a clear, easy justification without invoking anything about regular sequences; though it's certainly ok to say the word regular.

<sup>2</sup>Hint: show that any nonzero  $R$ -module homomorphism  $M \rightarrow R$  must be surjective, and then show that such a homomorphism must exist.

**Problem 6.** Let  $(R, \mathfrak{m})$  be a commutative local ring, and let  $M$  be a finitely presented  $R$ -module with minimal presentation

$$0 \longrightarrow K \longrightarrow R^n \xrightarrow{\pi} M \longrightarrow 0.$$

Note that the assumption here is that  $K$  is also a finitely generated module.

a) Show that if  $M$  is flat, then

$$0 \longrightarrow K \otimes_R R/\mathfrak{m} \longrightarrow R^n \otimes_R R/\mathfrak{m} \longrightarrow M \otimes_R R/\mathfrak{m} \longrightarrow 0$$

is exact.

b) Show that  $M$  is free  $\iff M$  is projective  $\iff M$  is flat.

**Problem 7.** ( $\star$ ) Let  $R$  be a domain and  $Q$  be its fraction field. Let  $T$  denote the torsion functor.

a) Show that  $T(M) = \text{Tor}_1^R(M, Q/R)$ .

b) Show that for every short exact sequence

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

of  $R$ -modules gives rise to an exact sequence

$$0 \longrightarrow T(A) \longrightarrow T(B) \longrightarrow T(C) \longrightarrow (Q/R) \otimes_R A \longrightarrow (Q/R) \otimes_R B \longrightarrow (Q/R) \otimes_R C \longrightarrow 0.$$

c) Show that the right derived functors of  $T$  are  $R^1T = (Q/R) \otimes_R -$  and  $R^iT = 0$  for all  $i \geq 2$ .

**Problem 8.** Let  $k$  be a field,  $R = k[x, y]$ , and  $\mathfrak{m} = (x, y)$ .

a) Show that

$$0 \longrightarrow R \xrightarrow{\begin{pmatrix} y \\ -x \end{pmatrix}} R^2 \xrightarrow{\begin{pmatrix} x & y \end{pmatrix}} R \longrightarrow 0$$

is a free resolution for  $k = R/\mathfrak{m}$ .

b) Compute  $\text{Tor}_i^R(k, k)$  for all  $i$ .

c) Show that

$$\text{Tor}_1(\mathfrak{m}, k) \cong \text{Tor}_2(k, k).$$