

Macaulay2 introductory problems

Problem 1.(a) Install Macaulay2.¹

Hardcore version: install emacs and run Macaulay2 through emacs.

(b) Make an .m2 file setting up a field k , a polynomial ring R over k , a nontrivial ideal I in R , the R -module $M = R/I$ and the ring $S = R/I$.**Problem 2** (Subalgebras). Use Macaulay2 to find:(a) A presentation for the \mathbb{Q} -algebra $\mathbb{Q}[xy, xu, yv, uv] \subseteq \mathbb{Q}[x, y, u, v]$.(b) A presentation for the k -algebra U , where $k = \mathbb{Z}/101$ and

$$k \begin{bmatrix} ux & uy & uz \\ vx & vy & vz \end{bmatrix} \subseteq \frac{k[u, v, x, y, z]}{(x^3 + y^3 + z^3)}.$$

Problem 3 (Graded rings).(a) In Macaulay2, set up $A = \mathbb{Q}[s^2, st, t^2]$ as an \mathbb{N}^2 -graded ring with the grading induced by setting s^2, st, t^2 as homogeneous elements of degrees

$$\deg(s^2) = (2, 0) \quad \deg(st) = (1, 1) \quad \deg(t^2) = (0, 2).$$

(b) The ring $R = \mathbb{Q}[t^3, t^{13}, t^{42}]$ is a graded subring of $\mathbb{Q}[t]$ with the standard grading, meaning that the graded structure on $\mathbb{Q}[t]$ induces a grading on R . Set up R (with this grading) in Macaulay2.**Problem 4** (Modules). Consider the domain $R = \mathbb{Q}[x, y, z, a, b, c]/(xb - ac, yc - bz, xc - az)$. Set up the following R -modules, making sure Macaulay2 actually sees them as modules over R :(a) The ideal $I = (x, a)$ viewed as an R -module.(b) The R -module $N = \mathbb{Q}$.(c) The 2-generated R -module $M = Rf + Rg$, where the generators f, g satisfy the relations

$$yf - xg = 0 \quad bf - cg = 0 \quad cf - zg = 0.$$

(d) The submodule of R^3 generated by (a, b, c) and (x, y, z) .**Problem 5** (Complexes in Macaulay2). Let $R = \mathbb{Q}[x, y, z]/(x^2, xy)$.

(a) Consider the bounded complex

$$C = \begin{array}{ccccccc} & & \begin{pmatrix} z \\ -y \\ x \end{pmatrix} & & \begin{pmatrix} -y & -z & 0 \\ x & 0 & -z \\ 0 & x & y \end{pmatrix} & & \begin{pmatrix} x & y & z \end{pmatrix} & & \\ & R & \longrightarrow & R^3 & \longrightarrow & R^3 & \longrightarrow & R & \\ & 3 & & 2 & & 1 & & 0 & \end{array}$$

Set C up in Macaulay2 and compute its homology. For which n is $H_n(C) = 0$?¹If your computer runs only Microsoft Windows, talk to Eloísa about it.

